

Receptor-ImmunoglobulinFc Chimeric Proteins Schematic Illustration




FIG. 1

Possible Nature of the LTB₄R-Ig
"Live/Dead" Problem




FIG. 2

3/11




FIG. 3




FIG. 4

09767370-061801

FACS ANALYSIS OF AG-H1-5 PURIFIED
UPPER AND LOWER BANDS ON PMA
(50ng/mL) ACTIVATED II-23 CELLS




FIG. 5A

FIG. 5B






FIG. 5C

5/11



TCBT90"02E29260





FIG. 6

6/11



FOBT90"02E29260




FIG. 7




FIG. 8

Effect of Incubation temperature
on % dead hLTBR




FIG. 9

09767370-061801

8/11





FIG. 10



9/11




FIG. 11C




FIG. 11B



37 C (4593 RU)
Ligand = human LIGHT




FIG. 12A

32 C (4760 RU)
Ligand = human LIGHT




FIG. 12B

28 C (4808 RU)
Ligand = human LIGHT




FIG. 12C

TOP SECRET 04E29260
10/11

37 C (4593 RU)
Ligand = human LTa




FIG. 12D

32 C (4760 RU)
Ligand = human LTa




FIG. 12E

28 C (4808 RU)
Ligand = human LTa




FIG. 12F